

AD-A154 160

MEMORY EFFECTS IN DYNAMICAL MANY-BODY SYSTEMS: THE  
ISOMNESIC (CONSTANT-ME. (U) ROCHESTER UNIV NY DEPT OF  
CHEMISTRY A C BERI ET AL. APR 85  
UROCHESTER/DC/85/TR-64 N00014-80-C-0472

1/1

UNCLASSIFIED


F/G 12/1

NL

END

FORMED

DTN



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

(2)

AD-A154 160

OFFICE OF NAVAL RESEARCH  
Contract N00014-80-C-0472  
Task No. NR 056-749  
TECHNICAL REPORT No. 64

Memory Effects in Dynamical Many-Body Systems:  
The Isomnesic (Constant-Memory) Approximation

by

A. C. Beri and Thomas F. George

Prepared for publication

in

Zeitschrift für Physik B: Condensed Matter

Department of Chemistry  
University of Rochester  
Rochester, New York 14627

April 1985

Reproduction in whole or in part is permitted for  
any purpose of the United States Government.

This document has been approved for public release  
and sale; its distribution is unlimited.

DTIC  
ELECTE  
MAY 29 1985  
S D  
B

DTIC FILE COPY

15 02 120

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UROCHESTER/DC/85/TR-64			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry University of Rochester		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research (Code 413)	
6c. ADDRESS (City, State and ZIP Code) River Station Rochester, New York 14627		7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-80-C-0472	
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO. 61153N		PROJECT NO. 013-08	TASK NO. NR 056-749
		WORK UNIT NO.			
11. TITLE Memory Effects in Dynamical Many-Body Systems: The Isomnesic (Constant-Memory) Approximation					
12. PERSONAL AUTHOR(S) A. C. Beri and Thomas F. George					
13a. TYPE OF REPORT Interim Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) April 1985	
15. PAGE COUNT 21					
16. SUPPLEMENTARY NOTATION Prepared for publication in Zeitschrift für Physik B: Condensed Matter					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	MANY-BODY SYSTEMS, GENERALIZED MASTER EQUATION, MEMORY EFFECTS, (continued on reverse side)		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) → Exact numerical solutions of a generalized master equation for a system involving a temporally localizable memory kernel and a constant one are presented. Comparison with an isomnesic (constant-memory) approximation, with the memory kernel reduced to the sum of a local and a constant term, gives excellent agreement. A Markovian version of the problem is also described and found to be inherently ambiguous, devoid of significant physical features seen in the transient regions, but leading to the same steady-state as the isomnesic approximation. → not required include:					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

18. ,ISOMNESIC APPROXIMATION,  
AGREEMENT WITH EXACT RESULTS,  
IMPROVEMENT ON MARKOVIAN APPROXIMATION.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

Memory Effects in Dynamical Many-Body Systems:  
The Isomnesic (Constant-Memory) Approximation

A. C. Beri and Thomas F. George  
Department of Chemistry  
University of Rochester  
Rochester, New York 14627

Abstract

Exact numerical solutions of a generalized master equation for a system involving a temporally localizable memory kernel and a constant one are presented. Comparison with an isomnesic (constant-memory) approximation, with the memory kernel reduced to the sum of a local and a constant term, gives excellent agreement. A Markovian version of the problem is also described and found to be inherently ambiguous, devoid of significant physical features seen in the transient regions, but leading to the same steady-state as the isomnesic approximation.

## I. Introduction

The dynamics of a system of many interacting particles is often treated by partitioning the system into two or more subsystems, one of which,  $\mathcal{S}$ , comprises the degree(s) of freedom of primary interest, while the others,  $\mathcal{R}_1, \mathcal{R}_2, \dots$ , are reservoirs to which  $\mathcal{S}$  is coupled.<sup>1</sup> No observations are made on the reservoirs, and the corresponding degrees of freedom are projected out of the equations of motion. A quantum mechanical generalized master equation (GME) results, and invocation of the Born and random-phase approximations generates a set of coupled Volterra-type integrodifferential equations (VTE)

$$\dot{P}_S(t) = \sum_{S' \neq S} \int_0^t dt' [K_{SS'}(t-t') P_{S'}(t') - K_{S'S}(t-t') P_S(t')] , \quad (1)$$

where  $P_S(t)$  is a diagonal matrix element of the projected density operator in the subspace  $\mathcal{S}$ , and represents the probability of finding  $\mathcal{S}$  in an eigenstate  $|S\rangle$  of the zero-order sub-Hamiltonian  $\mathcal{H}_S$ :

$$\mathcal{H}_S |S\rangle = \mathcal{E}_S |S\rangle . \quad (2)$$

The kernel functions  $K_{SS'}(t-t')$  include temporal correlations between dynamical variables of the reservoirs.<sup>2</sup> The resulting dependence of the state of  $\mathcal{S}$  at any time  $t$  upon all previous times  $t' \leq t$ , the memory effect, is manifested in the convolution form of  $K_{SS'}(t-t')$ , the memory kernels.<sup>3</sup> For most real physical systems, the form of  $K_{SS'}$  precludes an analytic closed-form solution, and often even step-by-step numerical integration.<sup>4</sup> When numerical solutions using alternative techniques such as the method of successive approximations<sup>5</sup> are possible, one is limited to small times. For example, for kernels with some form of periodicity, a fairly common situation, these times are of the order of

a few periods, which seldom correspond to macroscopic measurement times. The solutions are therefore of little use in suggesting or elucidating physical mechanisms.

The need for numerical solutions has, in the past, been circumvented by making the Markovian approximation (MA),<sup>6,7</sup> namely that the system is amnesic (has negligible memory). Thus the convolution-type kernels are assumed to be extremely localized temporally,

$$K_{SS}(t-t') = W_{SS} \delta(t-t'), \quad (3)$$

which is tantamount to assuming that the probability functions are very slowly varying during the (small) period of time in which  $K_{SS}(t)$  changes substantially. This approximation is difficult to justify in general, and is in fact not justified if a locally correct solution is sought. This is because  $P_S(t)$  must reflect the nature of  $K_{SS}(t)$  itself, and if the latter varies substantially in any time range, so must  $P_S(t)$ , at least locally in time. What makes the MA useful is its avowed ability to provide the rough behavior of  $P_S(t)$  over long time periods. Thus on a coarse-grained time scale, the MA is expected to provide results roughly resembling the exact results.

Not all physical systems admit of the representation of  $K_{SS}(t-t')$  shown in Eq. (3), an important example being a system resonantly pumped by a continuous-wave laser and coupled to another reservoir.<sup>7-9</sup> For this important class of physical systems the kernels can, under favorable conditions to be described later, be written as sums of terms such as Eq. (3) and additional constants independent of time.<sup>2</sup> The resulting equations of motion are very different from those obtained using Eq. (3). The differences are quantified and highlighted in this work, and it is shown that the results of the exactly-solvable equations with kernels including the constants provide an excellent



representation of the solutions. The MA, on the other hand, is shown to lead to probability profiles in gross disagreement with the exact results, especially for the short time scale.

The format of this paper is as follows. In Section II the basic theory is summarized, and the results of calculations are presented in Section III. Conclusions are given in Section IV.

## II. The Isomnesic (Constant-Memory) Approximation

A system of great current interest that lends itself ideally as a vehicle for illustrating the various approximations for the memory kernel is an adatom adsorbed on a solid surface, vibrationally excited by an infrared continuous-wave laser, and coupled to phonon modes of the solid.<sup>2,7-9</sup> The system  $S$  is the adbond, viz, the bond resulting from the interaction of the adatom with the phonon-averaged motion of the lattice atoms;<sup>10</sup> the phonons and the laser are two independent, noninteracting reservoirs  $\mathcal{R}_1$  and  $\mathcal{R}_2$ . Quantum theoretical treatments of this class of systems lead quite generally to a GME and a corresponding VTE, Eq. (1), with  $P_S(t)$  representing the probability of finding the adbond in the eigenstate  $|S\rangle$  of the zero-order adbond Hamiltonian  $\mathcal{H}_S$ . The kernels  $K_{SS'}(t-t')$  incorporate the influence of the phonons and the laser.<sup>2</sup> We have partitioned the total Hamiltonian of the system in such a way that the perturbations which generate the kernels are traceless in their individual reservoir subspaces.<sup>2,10</sup> As a direct consequence, the total kernels separate into individual subspace kernels,

$$K_{SS'}(t) = K_{SS'}^{(p)}(t) + K_{SS'}^{(r)}(t), \quad (4)$$

where  $K_{SS'}^{(p)}(t)$  is due to the phonon field alone and  $K_{SS'}^{(r)}(t)$  is due to the laser radiation field alone. Using a one-dimensional model for the solid,

we have obtained the kernels in closed form and solved them numerically by an iterative technique<sup>5</sup> for times of the order of a few "periods" of  $K_{SS'}(t)$ . These periods are dictated by the characteristic frequencies of the phonons and the adbond (i.e., the Debye frequency  $\omega_D$  and the transition frequencies  $\omega_{SS'}$  for bound states of the Hamiltonian  $\mathcal{H}_S$ ), and the laser frequency  $\omega_L$  all of which are  $\sim 10^{12} - 10^{14}$  Hz (1-100 THz). The periods are therefore very small, being  $\sim 10^{-12} - 10^{-14}$  s (1-.01 ps); consequently, the solutions of the VTE can be obtained numerically for times no larger than about 10 ps, which is certainly insufficient for the description of a physical or chemical process. In order to obtain solutions for longer times, an analytic solution must be sought.

We begin with the assertion that the physically important situation is one where the laser frequency and the Debye frequency equal different transition frequencies  $\omega_{SS'}$  of the adbond.<sup>2</sup> For this case  $K_{SS'}^{(r)}(t)$  becomes a constant  $k_{SS'}$  (from the term with  $|\omega_{SS'}| = \omega_L$ ) plus oscillating terms which we ignore in the sense of the rotating-wave approximation.<sup>6</sup> The phonon term,  $K_{SS'}^{(p)}(t)$ , has a complicated behavior typified by  $K_{01}^{(p)}(t)$  in Fig. 1, but we note that it has appreciable magnitude for times  $\sim 2$  ps, and almost vanishes subsequently. If the mesh size of our theoretical experiment is no less than 2 ps, we can treat  $K_{SS'}^{(p)}(t)$  as a delta function, so that, from Eq. (4),

$$K_{SS'}(t) = \Omega_{SS'} \delta(t) + k_{SS'} , \quad (5)$$

where

$$\Omega_{SS'} = \int_0^\infty dt K_{SS'}^{(p)}(t) . \quad (6)$$

In Eq. (5), the first term, originating from the coupling of the adbond to the phonons, is the amnesic or Markovian term, and would lead, in the absence of the laser term, to the elimination of all memory effects. The second term, due to

the resonant laser coupling, is the isomnesic non-Markovian term, and represents a constant memory effect all the way back to  $t=0$ .

The subsequent treatment is simplified by defining "diagonal" elements  $\Omega_{SS}$  and  $k_{SS}$ , viz.

$$\Omega_{SS} = - \sum_{S' \neq S} \Omega_{SS'}, \quad (7)$$

$$k_{SS} = - \sum_{S' \neq S} k_{S'S} \quad (8)$$

and the notation

$$\underline{\Omega} \equiv \{\Omega_{SS'}\}, \quad (9a)$$

$$\underline{k} \equiv \{k_{SS'}\}, \quad (9b)$$

$$\underline{k}(t) \equiv \{k_{SS'}(t)\}, \quad (9c)$$

$$\underline{w} \equiv \{w_{SS'}\}, \quad (9d)$$

$$\underline{\Delta} \equiv \{\delta_{SS'}\}, \quad (9e)$$

$$\underline{P}(t) \equiv \{P_S(t)\}. \quad (9f)$$

The exact VTE may be written as

$$\dot{\underline{P}}(t) = \int_0^t dt' \underline{k}(t-t') \cdot \underline{P}(t'); \quad (10)$$

in the isomnesic approximation (IA) it becomes

$$\dot{\underline{P}}(t) = \underline{k} \cdot \int_0^t dt' \underline{P}(t') + \underline{\Omega} \cdot \underline{P}(t). \quad (11)$$

Formally this can be solved without any further approximations. Differentiating both sides of Eq. (11), we get the second-order differential equation

$$\ddot{\underline{P}}(t) = \underline{k} \cdot \underline{P}(t) + \underline{\Omega} \cdot \dot{\underline{P}}(t), \quad (12)$$

whose Laplace transform,

$$\underline{M}(s) \cdot \underline{P}(s) = \underline{T}(s) , \quad (13)$$

where

$$\underline{M}(s) \equiv \underline{k} + s\underline{\Omega} - s^2 \underline{\Delta} , \quad (14)$$

$$\underline{T}(s) \equiv -s\underline{P}(0) , \quad (15)$$

and  $\underline{P}(s)$  is the Laplace transform of  $\underline{P}(t)$ , is easily inverted using the Heaviside expansion theorem.<sup>11</sup>

While this effectively eliminates the non-Markovian bottleneck without recourse to Eq. (3), it is instructive to generate solutions of a Markovian version of this class of problems. To do this, we have to make provisions for localizing  $K_{SS'}^{(r)}$  in time. This brings to the fore the pervasive issue of time scales in dynamical many-body systems, our specific question being the relative degree of localization ascribable to  $K_{SS'}^{(p)}$  and  $K_{SS'}^{(r)}$ . If the laser radiation is localized as a pulse, it can still extend over a period of time much longer than the extent of  $K_{SS'}^{(p)}(t)$ . In other words, the delta function representing  $K_{SS'}^{(p)}(t)$  may be quite different from the delta function representing  $K_{SS'}^{(r)}(t)$ , and the former may in fact not really be considered as localized with respect to the latter. With these provisos in mind, we can proceed to write the Markovian version of Eqs. (5), (10), (12) and (13):

$$\underline{K}(t) = (\underline{\Omega} + \underline{f}) \delta(t) , \quad (16)$$

$$\dot{\underline{P}}(t) = \underline{W} \cdot \underline{P}(t) , \quad (17)$$

$$\underline{J}(s) \cdot \underline{P}(s) = \underline{U} , \quad (18)$$

where

$$\underline{f} = \underline{k} \tau , \quad (19)$$

$$\underline{W} = \underline{\Omega} + \underline{f} , \quad (20)$$

$$\underline{J}(s) = (\underline{\Omega} + \underline{f} - s\underline{\Delta}) , \quad (21)$$

$$\underline{U} = -\underline{P}(0) , \quad (22)$$

and  $\tau$  represents the actual extent of the laser signal or a variable time parameter. Equation (17) is the well-known Pauli master equation (PME).<sup>1,6</sup>

To summarize the formulation, Eq. (10) is the exact GME with a convolution type kernel which makes solutions difficult for experimental times. If the kernel can be approximated as the sum of a very sharply peaked function due to the phonons plus a constant due to the laser (the isomnesic approximation), we get the simplified GME, Eq. (11), which is still not Markovian but has a "constant memory" (isomnesic-) term. The resulting second-order differential equation, Eq. (12), can be solved analytically for arbitrarily long times. If, on the other hand, memory effects are assumed to be negligible, i.e., with a kernel highly localized in time, Eq. (3), we get the Markovian limit, namely the PME, Eq. (17), which is a first-order differential equation and can also be solved analytically for arbitrarily long times. The nature of the solutions depends, of course, on the time parameter  $\tau$  in Eq. (19).

### III. Results of the Exact, Isomnesic and Markovian Approximations

For a linear-chain model of the solid, the exact kernel function due to phonons alone,  $K^{(p)}$ , involves an expansion over the number of phonons  $n$  and the number of lattice atoms  $l$ . For the specific system chosen, the effective surface potential generates five bound states; we have chosen  $|\omega_{01}| = \omega_D$  and  $|\omega_{13}| = \omega_L$  [ $\omega_{SS} = (\mathcal{E}_S - \mathcal{E}_S)/\hbar$ ; see Eq. (2)]. The phonon kernels, e.g.,  $K_{01}^{(p)}(t)$  in Fig. 1, fall off quite rapidly in time and arise almost entirely from the  $(n=1, l=1)$  term.

At issue is the validity of the isomnesic and Markovian approximations. We can examine two aspects of the results, viz. the details of the early transient behavior, and the long-term steady state. For the Markovian case, we can further examine the effects of varying  $\tau$  in Eq. (19). The transient behavior can be compared with exact numerical solutions of Eq. (10), but the steady-state results of the MA and the IA can only be judged by extrapolation.

While the probability profiles  $P(t)$  provide sufficient comparison, the physical contents of our results are better displayed by the average adbond energy given by

$$\mathcal{E}(t) = \sum_S P_S(t) \mathcal{E}_S, \quad (23)$$

where  $\mathcal{E}_S$  is the energy eigenvalue corresponding to the level  $|S\rangle$  of the effective adbond potential. This provides a simple measure of the chemical direction of the adbond and a well-defined connection between this exact theory and phenomenological models of such processes.

The results obtained by solving Eqs. (10), (12) and (17) for a variety of situations of physical interest are very consistent, and we present a typical set of curves in Fig. 2 for early times,  $t \leq 5$  ps. The exact results are rich in detail and, to a large degree, follow the variations in  $K(t)$ . We shall not dwell on the physical contents except to mention the conspicuous absence of the simple exponential approach to equilibrium postulated in relaxation time theories. The results of the IA, while exhibiting the expected loss of the fine detail, are seen to chart out the average behavior of  $P(t)$  with great accuracy. Thus, for example, the crossing point of  $P_1(t)$  and  $P_3(t)$  is reproduced at  $t = 1.7$  ps as compared to the exact crossing point at 2.2 ps, and the difference  $P_1(t) - P_3(t)$  is seen to follow the exact difference with a similar slight

temporal retardation. The gross oscillations of  $\underline{P}(t)$  are also reproduced faithfully. Consequently, we feel confident in our prediction that the steady-state results of the IA will conform to the exact results very accurately. We wish to emphasize the significance of this. The IA makes possible the complete solution of a GME for arbitrarily long times, when the solution of the exact GME is essentially impossible. The solution of the GME in the IA is no more difficult than that of the PME obtained by invoking the MA which, however, gives results which differ radically from the exact results in the transient region for a wide range of values of  $\tau$ . Figure 3 illustrates the variations of  $P_S(t)$  with  $\tau$  in the MA ( $10^{-5}$  ps  $\leq \tau \leq 10^5$  ps). The probability profiles are seen to be insensitive to increases in  $\tau$  beyond  $\sim 10$  ps, which corresponds to almost instantaneous saturation. The apparent insensitivity of  $\underline{P}(t)$  for  $\tau \leq 10^{-2}$  ps seen in Fig. 3 is, of course, misleading, since in this region the  $P_S(t)$  vary with time too slowly for the change to be evident in the 20 ps time range of the plot. Drawn for 2 ns in Fig. 4 [we present only  $P_3(t)$ ], the differences become more pronounced for  $\tau \leq 10^{-2}$  ps, and the strong dependence of the apparent "equilibration time" on  $\tau$  emerges clearly. We see no unambiguous way of choosing an appropriate value for  $\tau$ . For systems which are known or expected to equilibrate very quickly, one could make an ad hoc semiempirical choice of  $\tau$  to solve a PME for the steady state. However, if one is interested in the details of the short-time behavior of the system, one must solve the GME using the IA.

For comparison of the MA with the IA we have chosen  $\tau = 1$  ps. The resulting profiles for the MA are shown in Fig. 5, along with our IA results for 20 ps. The complete absence of oscillatory behavior representing beats of the system and of population crossings, and the presence of predominantly exponential variation of  $\underline{P}(t)$  representative of a "relaxation time" exhibited by the MA

results of Fig. 5 are not necessarily general features of the solutions of a PME. Since the matrices  $\underline{M}$  and  $\underline{J}$  are, in general, asymmetric, both the GME and the PME can generate oscillatory solutions  $\underline{P}(t)$ , contradicting earlier contentions.<sup>12</sup> However, the total lack of resemblance between the early PME results and the exact ones for the present case is alarming and raises urgent and serious questions about the validity of the Markovian approximation, even for a rough qualitative estimate of the early transient behavior of many-body systems with a nonrandom force. At best, these results suggest the need for great caution in using the MA. In retrospect this is not surprising, because the representation of a time-independent cw laser beam by an ultrashort pulse is tenuous, whereas that of the very-short-range (in time) phonon correlation by a delta function is quite reasonable.

The results for  $\mathcal{E}(t)$ , shown in Fig. 6 for the exact GME and the IA and in Fig. 7 for the MA and the IA further support these contentions. The IA generates an energy profile which, except for a slight temporal retardation, is a low-resolution representation of the exact result. The significant increase in the bond energy at the expense of both the laser and the solid is clearly evident in the IA results of Figs. 2 and 6. The results of the MA are qualitatively different from the exact and IA results, and describe a monotonic approach to the steady state. The exact and IA results show actual periodic energy transfer out of the adbond followed by successive build up due to overall energy inflow. The apparent correspondence seen in Fig. 7 between the magnitude of  $\mathcal{E}(t)$  in the MA and the "base line" about which the results of the exact GME and the IA oscillate is entirely due to our choice of  $\tau = 1$  ps. That no such correspondence can be expected in general is evident from Fig. 8 showing the variations of  $\mathcal{E}(t)$  with  $\tau$ . The steady-state ( $t \rightarrow \infty$ ) values of  $\underline{P}(t)$  and  $\mathcal{E}(t)$  in



the MA have been found to be the same as those obtained with the IA for all cases considered so far (when the values of  $\tau$  chosen lead to steady states in the time ranges considered).

#### IV. Conclusion

Our results for the time evolution of a dynamical many-body system subject to a variety of forces point out the inadequacies of the Markovian approximation when the resulting memory kernel is nonlocal in time. As an alternative, we present the isomnesic approximation which is characterized by a constant memory; this eliminates the non-Markovian bottleneck associated with the convolution-type memory kernel in the Volterra-type integrodifferential equation resulting from the generalized master equation. Closed solutions of the latter are in excellent agreement with the exact results, in contrast to those of the Markovian approximation.

#### Acknowledgements

This research was supported by the Office of Naval Research, the Air Force Office of Scientific Research (AFSC), United States Air Force under Grant AFOSR-82-0046, and the National Science Foundation under Grant CHE-8320185. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. TFG acknowledges the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Award (1975-86).

### References

1. F. Haake, in Springer Tracts in Modern Physics, edited by G. Höhler (Springer-Verlag, Berlin, 1973), p. 98ff.
2. A. C. Beri and Thomas F. George, unpublished.
3. R. W. Zwanzig, in Lectures in Theoretical Physics, Vol. III, edited by W. E. Brittin, B. W. Downs and J. Downs (Interscience Publishers, New York, 1961), p. 106ff.
4. See, for example, Treatment of Integral Equations by Numerical Methods edited by C. T. H. Baker and G. F. Miller (Academic Press, New York, 1982) and Numerical Treatment of Integral Equations edited by J. Albrecht, Clausthal-Zellerfeld and L. Collatz (Birkhäuser Verlag, Basel, 1980).
5. L. Collatz, "The Numerical Treatment of Differential Equations" (Springer-Verlag, New York, 1966), Chapter VI.
6. W. H. Louisell, "Quantum Statistical Properties of Radiation" (John Wiley & Sons, New York, 1973), Chapter 6.
7. J. Lin, A. C. Beri, M. Hutchinson, W. C. Murphy and T. F. George, Phys. Lett. 79A, 233 (1980).
8. T. F. George, A. C. Beri, K. S. Lam and J. Lin, in Laser Applications, edited by J. F. Ready and R. K. Erf (Academic Press, New York, 1984), pp. 69-127; T. F. George, J. Lin, A. C. Beri and W. C. Murphy, Prog. Surf. Sci., in press.
9. C. Jedrzejek, K. F. Freed, S. Efrima and H. Metiu, Surf. Sci. 109, 191 (1981).
10. C. Jedrzejek, K. F. Freed, E. Hood and H. Metiu, J. Chem. Phys. 79, 2436 (1983).
11. E. Butkov, "Mathematical Physics", (Addison Wesley Publishing Co., Reading, Massachusetts, 1968), Chapter 5.
12. A. Janner, L. Van Hove and E. Verboven, Physica 28, 1341 (1962).

### Figure Captions

Fig. 1. The phonon part  $K_{01}^{(P)}(t)$  of the memory kernel.

Fig. 2. Probability profiles  $P_S(t)$  for levels  $S=0, 1$  and  $3$  of the adbond. Solid lines: Exact numerical solutions of the GME; broken lines: closed-form solutions of the GME with the IA.

Fig. 3.  $P_S(t)$  for the system of Fig. 2 with the MA for a ten-decade range of  $\tau$ . (A)- $P_0(t)$ ; (B)- $P_1(t)$ ; (C)- $P_3(t)$ .

Fig. 4.  $P_3(t)$  with the MA for 2 ns.


Fig. 5. Comparison of  $P_S(t)$  with the IA (solid lines) and the MA ( $\tau=1$  ps; broken lines) for  $S=0, 1$  and  $3$ .

Fig. 6. Average adbond energy  $\mathcal{E}(t)$  obtained via the numerical solution of the exact GME (solid line) and closed-form solution with the IA (broken line).

Fig. 7.  $\mathcal{E}(t)$  with the IA (solid line) and the MA ( $\tau=1$  ps; broken line).


Fig. 8. Variation of  $\mathcal{E}(t)$  with  $\tau$  in the MA.

## MEMORY KERNEL




Ber/Georgas Fig 1

## EXACT vs. ISOMNESIC



Ber/Georgas Tr 2


# MARKOVIAN APPROXIMATION



(A)

Bevi/George Fig 3(A)


# MARKOVIAN APPROXIMATION



(B)


Bevi/George Fig 3(B)

# MARKOVIAN APPROXIMATION




Beri / George Fig 3(c)

# MARKOVIAN APPROXIMATION




Beri / George Fig 4

## *ISOMNESIC vs. MARKOVIAN*




Beri/George Fig 5

## *EXACT vs. ISOMNESIC*




Beri/George Fig 6

# ISOMNESIC vs. MARKOVIAN



Ber / George Fig 7

## MARKOVIAN APPROXIMATION



Ber / George Fig 8



DL/413/83/01  
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Doua Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Surface Chemistry Division (6170)  
455 Overlook Avenue, S.W.  
Washington, D.C. 20375

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. P. Lund  
Department of Chemistry  
Howard University  
Washington, D.C. 20059

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6132  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Richard Colton  
Code 6112  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 5570  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
California Institute of Technology  
Division of Chemistry and Chemical  
Engineering  
Pasadena, California 91125

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
Southampton University  
Southampton SO9 5NH  
Hampshire, England

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

Dr. Richard Smardzewski  
Code 6130  
Naval Research Laboratory  
Washington, D.C. 20375

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

DL/413/83/01  
056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

~~Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627~~

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Captain Lee Myers  
AFOSR/NC  
Bollig AFB  
Washington, D.C. 20332

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 41106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. G. D. Stein  
Mechanical Engineering Department  
Northwestern University  
Evanston, Illinois 60201

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G. H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. P. Hansma  
Physics Department  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
California Institute of Technology  
Division of Chemistry  
Pasadena, California 91125

Dr. W. Goddard  
California Institute of Technology  
Division of Chemistry  
Pasadena, California 91125

DL/413/83/01  
056/413-2

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

**END**

**FILMED**

**7-85**

**DTIC**